arXiv:1304.1591v1 [quant-ph] 5 Apr 2013

"Proof" of the Copenhagen Interpretation

Kazuyuki FUJII [*]

International College of Arts and Sciences

Yokohama City University

Yokohama, 236–0027

Japan

Abstract

When we start studying Quantum Mechanics the most difficult part to understand is the so–called Copenhagen interpretation. Usually beginners skip over this part, which is a wise choice in a certain sense. However, some researchers feel guilty about skipping over this.

In this paper we try to give a proof to it. Our method is based on embedding it into decoherence theory. The method is of course not complete, but some researchers may feel relieved.

To the best of our knowledge this is the finest method at the present time.

Keywords : quantum mechanics; Copenhagen interpretation; decoherence theory.

1 Introduction

Let us start with a system of principles of Quantum Mechanics (QM for simplicity). See for example [1], [2] and [3]. That is,

---

[*] E-mail address : fujii@yokohama-cu.ac.jp

System of Principles of QM

1. Superposition Principle

If $|a\rangle$ and $|b\rangle$ are physical states then superposition $\alpha|a\rangle + \beta|b\rangle$ is also a physical state where α and β are complex numbers.

2. Schrödinger Equation and Evolution

Time evolution of a physical state proceeds like

$$|\Psi\rangle \longrightarrow U(t)|\Psi\rangle$$

where $U(t)$ is the unitary evolution operator ($U^\dagger(t)U(t) = U(t)U^\dagger(t) = \mathbf{1}$ and $U(0) = \mathbf{1}$) determined by a Schrödinger Equation.

3. Copenhagen Interpretation[1]

Let a and b be eigenvalues of an observable Q, and $|a\rangle$ and $|b\rangle$ be eigenstates corresponding to a and b. When a state is superposition $\alpha|a\rangle + \beta|b\rangle$ and we observe the observable Q the state collapses like

$$\alpha|a\rangle + \beta|b\rangle \longrightarrow |a\rangle \quad \text{or} \quad \alpha|a\rangle + \beta|b\rangle \longrightarrow |b\rangle$$

where their collapsing probabilities are $|\alpha|^2$ and $|\beta|^2$ respectively ($|\alpha|^2 + |\beta|^2 = 1$).

This is called the collapse of wave function and probabilistic interpretation.

4. Many Particle State and Tensor Product

A multiparticle state becomes superposition of the Knonecker product of one particle states, which is called the tensor product. For example,

$$\alpha|a\rangle \otimes |a\rangle + \beta|b\rangle \otimes |b\rangle \equiv \alpha|a, a\rangle + \beta|b, b\rangle$$

is a two particle state.

---

[1] There are some researchers who are against this terminology, see for example [3]. However, I don't agree with them because the terminology is nowadays very popular in the world

The target of this paper is to give a proof to the Copenhagen Interpretation, so we give a symbolic figure of it for the latter convenience (we take $|0\rangle$ and $|1\rangle$ in place of $|a\rangle$ and $|b\rangle$ in the following).



$|0\rangle$
$\alpha|0\rangle+\beta|1\rangle$
Detector
$|0\rangle$ (probability $|\alpha|^2$)
$\alpha|0\rangle+\beta|1\rangle$
$|1\rangle$ (probability $|\beta|^2$)
time
$t=0$
$t=t_0$

Figure : Image of the Copenhagen Interpretation

2 "Proof" of the Copenhagen Interpretation

In this section we try to give a proof to the Copenhagen Interpretation. We perform this by embedding it into decoherence theory.

We consider an atom flying as in the figure of the preceding section and treat a two level system of the atom in the following, see for example [4]. First of all let us prepare some notations from Quantum Optics. Since we treat the two level system of the atom the target space is $\mathbf{C}^2 = \mathrm{Vect}_{\mathbf{C}}(|0\rangle, |1\rangle)$ with bases

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Then Pauli matrices $\{\sigma_1,\ \sigma_2,\ \sigma_3\}$ with the identity 1_2

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad 1_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

act on the space. For

$$\sigma_+ \equiv \frac{1}{2}(\sigma_1 + i\sigma_2) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \sigma_- \equiv \frac{1}{2}(\sigma_1 - i\sigma_2) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

it is easy to see

$$\sigma_+\sigma_- = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \sigma_-\sigma_+ = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Here we may assume that the initial state is $|0\rangle$ at $t = 0$ and the intermediate state is $\alpha|0\rangle + \beta|1\rangle$ for $0 < t < t_0$ and the last state is the one detected at $t = t_0$, see the figure in the preceding section once more.

For the initial time $t = 0$ we can assume that the Hamiltonian is a diagonal form

$$H_0 = \begin{pmatrix} E_0 & 0 \\ 0 & E_1 \end{pmatrix} \tag{1}$$

where E_0 and E_1 are the eigenvalues ($E_0 < E_1$ for simplicity) of the atom. It is easy to see

$$H_0|0\rangle = E_0|0\rangle, \quad H_0|1\rangle = E_1|1\rangle.$$

For the intermediate time $0 < t < t_0$ the Hamiltonian becomes

$$\begin{aligned} H &= U(\alpha, \beta) \begin{pmatrix} E_0 & 0 \\ 0 & E_1 \end{pmatrix} U(\alpha, \beta)^\dagger \\ &= \begin{pmatrix} |\alpha|^2 E_0 + |\beta|^2 E_1 & \alpha\bar{\beta}(E_0 - E_1) \\ \bar{\alpha}\beta(E_0 - E_1) & |\beta|^2 E_0 + |\alpha|^2 E_1 \end{pmatrix} \end{aligned} \tag{2}$$

where $U(\alpha,\beta)$ is a (special) unitary matrix given by

$$U = U(\alpha,\beta) = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} \quad (|\alpha|^2 + |\beta|^2 = 1). \tag{3}$$

In this case, it is easy to see

$$\alpha|0\rangle + \beta|1\rangle = \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \quad \text{and} \quad H(\alpha|0\rangle + \beta|1\rangle) = E_0(\alpha|0\rangle + \beta|1\rangle).$$

Note that H and H_0 are of course hermitian matrices ($H = H^\dagger$, $H_0 = H_0^\dagger$).

Note Since

$$(\alpha|0\rangle + \beta|1\rangle)(\alpha|0\rangle + \beta|1\rangle)^\dagger = |\alpha|^2|0\rangle\langle 0| + \alpha\bar{\beta}|0\rangle\langle 1| + \bar{\alpha}\beta|1\rangle\langle 0| + |\beta|^2|1\rangle\langle 1|$$

the Copenhagen interpretation may be written as collapsing

$$(\alpha|0\rangle + \beta|1\rangle)(\alpha|0\rangle + \beta|1\rangle)^\dagger \longrightarrow |\alpha|^2|0\rangle\langle 0| + |\beta|^2|1\rangle\langle 1|.$$

The method developped in the following is based on the paper [5]. In order to embed the Copenhagen Interpretation into decoherence theory we consider hypothetical time T and take it to be $T = t/(t - t_0)$ for example. Then

$$t = 0 \Longrightarrow T = 0\ ; \quad t = t_0 \Longrightarrow T = \infty. \tag{4}$$

However, for simplicity we again set $T = t$ in the following (we believe that there is no confusion).

To treat decoherence in a correct manner we must change models based on from a pure state to a density matrix. The general definition of density matrix ρ is given by both $\rho^\dagger = \rho$ and $\mathrm{tr}\rho = 1$, so we can write $\rho = \rho(t)$ as

$$\rho = \begin{pmatrix} a & b \\ \bar{b} & d \end{pmatrix} \quad (a = \bar{a},\ d = \bar{d},\ a + d = 1). \tag{5}$$

The general form of master equation ([6], [7]) is well–known to be

$$\frac{d}{dt}\rho = -i[H,\rho] + D\rho \quad (\Leftarrow \hbar = 1 \text{ for simplicity}) \tag{6}$$

where

$$D\rho=\mu\left(\sigma_-\rho\sigma_+-\frac{1}{2}\sigma_+\sigma_-\rho-\frac{1}{2}\rho\sigma_+\sigma_-\right)+\nu\left(\sigma_+\rho\sigma_--\frac{1}{2}\sigma_-\sigma_+\rho-\frac{1}{2}\rho\sigma_-\sigma_+\right)$$

and $\mu,\ \nu>0$. Note that μ and ν are important constants determined later.

We must solve the equation. If we write H in (2) as

$$H=\begin{pmatrix} h & k \\ \bar{k} & l \end{pmatrix} \quad (h,\ l\in\mathbf{R},\ k\in\mathbf{C}) \tag{7}$$

for simplicity, then the master equation above can be rewritten as

$$\frac{d}{dt}\begin{pmatrix} a \\ b \\ \bar{b} \\ d \end{pmatrix}=\begin{pmatrix} -\mu & i\bar{k} & -ik & \nu \\ ik & i(l-h)-\frac{\mu+\nu}{2} & 0 & -ik \\ -i\bar{k} & 0 & -i(l-h)-\frac{\mu+\nu}{2} & i\bar{k} \\ \mu & -i\bar{k} & ik & -\nu \end{pmatrix}\begin{pmatrix} a \\ b \\ \bar{b} \\ d \end{pmatrix}. \tag{8}$$

We leave the derivation to readers.

Note and set

$$\begin{aligned}
&\begin{pmatrix} -\mu & i\bar{k} & -ik & \nu \\ ik & i(l-h)-\frac{\mu+\nu}{2} & 0 & -ik \\ -i\bar{k} & 0 & -i(l-h)-\frac{\mu+\nu}{2} & i\bar{k} \\ \mu & -i\bar{k} & ik & -\nu \end{pmatrix} \\
=&\begin{pmatrix} 0 & i\bar{k} & -ik & 0 \\ ik & i(l-h) & 0 & -ik \\ -i\bar{k} & 0 & -i(l-h) & i\bar{k} \\ 0 & -i\bar{k} & ik & 0 \end{pmatrix}+\begin{pmatrix} -\mu & 0 & 0 & \nu \\ 0 & -\frac{\mu+\nu}{2} & 0 & 0 \\ 0 & 0 & -\frac{\mu+\nu}{2} & 0 \\ \mu & 0 & 0 & -\nu \end{pmatrix} \\
\equiv&\ \widehat{H}+\widehat{D}.
\end{aligned}$$

The general solution of (8) is given by

$$\begin{pmatrix} a(t) \\ b(t) \\ \bar{b}(t) \\ d(t) \end{pmatrix}=e^{t\left(\widehat{H}+\widehat{D}\right)}\begin{pmatrix} a(0) \\ b(0) \\ \bar{b}(0) \\ d(0) \end{pmatrix}. \tag{9}$$

However, it is not easy to calculate the term $e^{t\left(\widehat{H}+\widehat{D}\right)}$ exactly, so we use a simple approximation

$$e^{t\left(\widehat{H}+\widehat{D}\right)}=e^{t\left(\widehat{D}+\widehat{H}\right)}\approx e^{t\widehat{D}}e^{t\widehat{H}}.$$

In general, we must use the Zassenhaus formula (see for example [8], [9]).

Zassenhaus Formula For operators (or square matrices) A and B we have an expansion

$$e^{t(A+B)}=\cdots e^{-\frac{t^3}{6}\{2[[A,B],B]+[[A,B],A]\}}e^{\frac{t^2}{2}[A,B]}e^{tB}e^{tA}. \tag{10}$$

The formula is a bit different from that of [8].

Therefore, we treat the approximate solution

$$\begin{pmatrix} a(t) \\ b(t) \\ \bar{b}(t) \\ d(t) \end{pmatrix}\approx e^{t\widehat{D}}e^{t\widehat{H}}\begin{pmatrix} a(0) \\ b(0) \\ \bar{b}(0) \\ d(0) \end{pmatrix}. \tag{11}$$

First, let us calculate $e^{t\widehat{D}}$. For the purpose we set

$$K=\begin{pmatrix} -\mu & \nu \\ \mu & -\nu \end{pmatrix}$$

and calculate e^{tK}. The eigenvalues of K are $\{0,-(\mu+\nu)\}$ and corresponding eigenvectors (not normalized) are

$$0\longleftrightarrow\begin{pmatrix} \nu \\ \mu \end{pmatrix},\quad -(\mu+\nu)\longleftrightarrow\begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

If we define the matrix

$$O=\begin{pmatrix} \nu & 1 \\ \mu & -1 \end{pmatrix}\Longrightarrow O^{-1}=\frac{1}{\mu+\nu}\begin{pmatrix} 1 & 1 \\ \mu & -\nu \end{pmatrix}$$

then it is easy to see

$$K=O\begin{pmatrix} 0 & \\ & -(\mu+\nu) \end{pmatrix}O^{-1}$$

and

$$e^{tK}=O\begin{pmatrix}1 & \\ & e^{-t(\mu+\nu)}\end{pmatrix}O^{-1}=\frac{1}{\mu+\nu}\begin{pmatrix}\nu+\mu e^{-t(\mu+\nu)} & \nu-\nu e^{-t(\mu+\nu)}\\ \mu-\mu e^{-t(\mu+\nu)} & \mu+\nu e^{-t(\mu+\nu)}\end{pmatrix}.$$

Therefore, we have

$$e^{t\widehat{D}}=\begin{pmatrix}\frac{\nu+\mu e^{-t(\mu+\nu)}}{\mu+\nu} & 0 & 0 & \frac{\nu-\nu e^{-t(\mu+\nu)}}{\mu+\nu}\\ 0 & e^{-t\frac{\mu+\nu}{2}} & 0 & 0\\ 0 & 0 & e^{-t\frac{\mu+\nu}{2}} & 0\\ \frac{\mu-\mu e^{-t(\mu+\nu)}}{\mu+\nu} & 0 & 0 & \frac{\mu+\nu e^{-t(\mu+\nu)}}{\mu+\nu}\end{pmatrix}\approx\frac{1}{\mu+\nu}\begin{pmatrix}\nu & 0 & 0 & \nu\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0\\ \mu & 0 & 0 & \mu\end{pmatrix} \qquad (12)$$

if t is large enough ($t \gg 1/(\mu+\nu)$).

Next, let us calculate $e^{t\widehat{H}}$. Since we need some properties of tensor product in the following see for example [9]. We can write the equation as

$$\widehat{H}=-i\left(H\otimes 1_2-1_2\otimes H^T\right).$$

In fact,

$$\begin{aligned}\widehat{H} &= -i\left\{\begin{pmatrix}h & k\\ \bar{k} & l\end{pmatrix}\otimes\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}-\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}\otimes\begin{pmatrix}h & \bar{k}\\ k & l\end{pmatrix}\right\}\\ &= -i\left\{\begin{pmatrix}h & 0 & k & 0\\ 0 & h & 0 & k\\ \bar{k} & 0 & l & 0\\ 0 & \bar{k} & 0 & l\end{pmatrix}-\begin{pmatrix}h & \bar{k} & 0 & 0\\ k & l & 0 & 0\\ 0 & 0 & h & \bar{k}\\ 0 & 0 & k & l\end{pmatrix}\right\}\\ &= -i\begin{pmatrix}0 & -\bar{k} & k & 0\\ -k & -(l-h) & 0 & k\\ \bar{k} & 0 & l-h & -\bar{k}\\ 0 & \bar{k} & -k & 0\end{pmatrix}.\end{aligned}$$

It is well–known that

$$e^{t\widehat{H}}=e^{-it\left(H\otimes 1_2-1_2\otimes H^T\right)}=e^{-itH\otimes 1_2}e^{it1_2\otimes H^T}=\left(e^{-itH}\otimes 1_2\right)\left(1_2\otimes e^{itH^T}\right)=e^{-itH}\otimes e^{itH^T}.$$

Since

$$H = U \begin{pmatrix} E_0 & 0 \\ 0 & E_1 \end{pmatrix} U^\dagger$$

we have

$$e^{-itH} = U \begin{pmatrix} e^{-itE_0} & 0 \\ 0 & e^{-itE_1} \end{pmatrix} U^\dagger$$

and

$$\begin{aligned} e^{t\widehat{H}} &= \left\{ U \begin{pmatrix} e^{-itE_0} & 0 \\ 0 & e^{-itE_1} \end{pmatrix} U^\dagger \right\} \otimes \left\{ U \begin{pmatrix} e^{itE_0} & 0 \\ 0 & e^{itE_1} \end{pmatrix} U^\dagger \right\}^T \\ &= U \otimes (U^\dagger)^T \left\{ \begin{pmatrix} e^{-itE_0} & 0 \\ 0 & e^{-itE_1} \end{pmatrix} \otimes \begin{pmatrix} e^{itE_0} & 0 \\ 0 & e^{itE_1} \end{pmatrix} \right\} \left(U \otimes (U^\dagger)^T \right)^\dagger \\ &= U \otimes (U^\dagger)^T \begin{pmatrix} 1 & & & \\ & e^{it(E_1 - E_0)} & & \\ & & e^{-it(E_1 - E_0)} & \\ & & & 1 \end{pmatrix} \left(U \otimes (U^\dagger)^T \right)^\dagger . \end{aligned}$$

Here we have used well–known formulas on tensor product

$$(A_1 \otimes B_1)(A_2 \otimes B_2) = A_1 A_2 \otimes B_1 B_2,\ (A \otimes B)^\dagger = A^\dagger \otimes B^\dagger,\ (A \otimes B)^T = A^T \otimes B^T,$$

see for example [9].

Since

$$U = U(\alpha, \beta) = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} \quad (|\alpha|^2 + |\beta|^2 = 1)$$

from (3) we have

$$U \otimes (U^\dagger)^T = \begin{pmatrix} |\alpha|^2 & -\alpha\beta & -\bar{\alpha}\bar{\beta} & |\beta|^2 \\ \alpha\bar{\beta} & \alpha^2 & -\bar{\beta}^2 & -\alpha\bar{\beta} \\ \bar{\alpha}\beta & -\beta^2 & \bar{\alpha}^2 & -\bar{\alpha}\beta \\ |\beta|^2 & \alpha\beta & \bar{\alpha}\bar{\beta} & |\alpha|^2 \end{pmatrix}$$

and, by setting $J = e^{it(E_1 - E_0)}$ for simplicity,

$$\begin{aligned}
e^{t\widehat{H}} &= U \otimes (U^\dagger)^T \begin{pmatrix} 1 & & & \\ & J & & \\ & & J^{-1} & \\ & & & 1 \end{pmatrix} \left(U \otimes (U^\dagger)^T \right)^\dagger \\
&= \begin{pmatrix} c_{11} & c_{12} & c_{13} & c_{14} \\ * & * & * & * \\ * & * & * & * \\ c_{41} & c_{42} & c_{43} & c_{44} \end{pmatrix} \qquad (13)
\end{aligned}$$

where

$$\begin{aligned}
c_{11} &= |\alpha|^4 + (J + J^{-1})|\alpha|^2|\beta|^2 + |\beta|^4, \\
c_{12} &= \left(|\alpha|^2 - |\alpha|^2 J + |\beta|^2 J^{-1} - |\beta|^2 \right) \bar{\alpha}\beta, \\
c_{13} &= \left(|\alpha|^2 + |\beta|^2 J - |\alpha|^2 J^{-1} - |\beta|^2 \right) \alpha\bar{\beta}, \\
c_{14} &= (2 - J - J^{-1})|\alpha|^2|\beta|^2
\end{aligned}$$

and

$$\begin{aligned}
c_{41} &= (2 - J - J^{-1})|\alpha|^2|\beta|^2, \\
c_{42} &= \left(|\beta|^2 + |\alpha|^2 J - |\beta|^2 J^{-1} - |\alpha|^2 \right) \bar{\alpha}\beta, \\
c_{43} &= \left(|\beta|^2 - |\beta|^2 J + |\alpha|^2 J^{-1} - |\alpha|^2 \right) \alpha\bar{\beta}, \\
c_{44} &= |\beta|^4 + (J + J^{-1})|\alpha|^2|\beta|^2 + |\alpha|^4.
\end{aligned}$$

Note that $*$'s in the matrix are elements not used in the latter. We leave the derivation to readers.

Here, we list very important relations among $\{\alpha\}$ (coming from $|\alpha|^2 + |\beta|^2 = 1$)

$$c_{11} + c_{41} = 1, \quad c_{12} + c_{42} = 0, \quad c_{13} + c_{43} = 0, \quad c_{14} + c_{44} = 1. \qquad (14)$$

Therefore, from (11), (12), (13) and (14) we obtain

$$\begin{aligned}\begin{pmatrix} a(t) \\ b(t) \\ \bar{b}(t) \\ d(t) \end{pmatrix} &\approx \frac{1}{\mu+\nu}\begin{pmatrix} \nu & 0 & 0 & \nu \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \mu & 0 & 0 & \mu \end{pmatrix}\begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} \\ * & * & * & * \\ * & * & * & * \\ \alpha_{41} & \alpha_{42} & \alpha_{43} & \alpha_{44} \end{pmatrix}\begin{pmatrix} a(0) \\ b(0) \\ \bar{b}(0) \\ d(0) \end{pmatrix} \\ &= \frac{1}{\mu+\nu}\begin{pmatrix} \nu & 0 & 0 & \nu \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \mu & 0 & 0 & \mu \end{pmatrix}\begin{pmatrix} a(0) \\ b(0) \\ \bar{b}(0) \\ d(0) \end{pmatrix} \end{aligned} \tag{15}$$

for $t \gg 1/(\mu+\nu)$.

From (5)

$$\rho(t) = \begin{pmatrix} a(t) & b(t) \\ \bar{b}(t) & d(t) \end{pmatrix}$$

we have

$$\rho(\infty) = \frac{1}{\mu+\nu}\begin{pmatrix} \nu\,(a(0)+d(0)) & 0 \\ 0 & \mu\,(a(0)+d(0)) \end{pmatrix}.$$

The initial density matrix

$$\rho(0) = |0\rangle\langle 0| = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \equiv \begin{pmatrix} a(0) & b(0) \\ \bar{b}(0) & d(0) \end{pmatrix}$$

gives

$$\rho(\infty) = \frac{1}{\mu+\nu}\begin{pmatrix} \nu & 0 \\ 0 & \mu \end{pmatrix} = \frac{\nu}{\mu+\nu}|0\rangle\langle 0| + \frac{\mu}{\mu+\nu}|1\rangle\langle 1|. \tag{16}$$

Here, if we choose

$$\mu = |\beta|^2 \quad \text{and} \quad \nu = |\alpha|^2 \quad (|\alpha|^2 + |\beta|^2 = 1) \tag{17}$$

from the starting point then we have the final form

$$\rho(\infty) = |\alpha|^2|0\rangle\langle 0| + |\beta|^2|1\rangle\langle 1|. \tag{18}$$

We can interpret this equation as a mathematical expression of the Copenhagen interpretation : "when a state is superposition $\alpha|0\rangle + \beta|1\rangle$ and we observe the observable Q the state collapses like $\alpha|a\rangle+\beta|b\rangle \ \rightarrow\ |a\rangle$ (probability $|\alpha|^2$) or $\alpha|a\rangle+\beta|b\rangle \ \rightarrow\ |b\rangle$ (probability $|\beta|^2$)."

This finishes the "proof" of the Copenhagen Interpretation.

Problem Our calculation is based on a simple approximation, so give a full calculation.

3 Summary

In this section let us summary our method briefly for the sake of readers.

[A] $t = 0$:

Hamiltonian H_0

$$H_0|0\rangle = E_0|0\rangle, \quad H_0|1\rangle = E_1|1\rangle.$$

Density matrix $\rho(0) = |0\rangle\langle 0|$.

[B] $t > 0$:

Hamiltonian $H = U(\alpha,\beta)H_0U(\alpha,\beta)^\dagger$ $(|\alpha|^2 + |\beta|^2 = 1)$

$$H(\alpha|0\rangle + \beta|1\rangle) = E_0(\alpha|0\rangle + \beta|1\rangle), \quad H(-\bar{\beta}|0\rangle + \bar{\alpha}|1\rangle) = E_1(-\bar{\beta}|0\rangle + \bar{\alpha}|1\rangle).$$

Density matrix $\rho(t)$. This is determined by the (special) master equation

$$\frac{d}{dt}\rho = -i[H,\rho]+|\beta|^2\left(\sigma_-\rho\sigma_+ - \frac{1}{2}\sigma_+\sigma_-\rho - \frac{1}{2}\rho\sigma_+\sigma_-\right)+|\alpha|^2\left(\sigma_+\rho\sigma_- - \frac{1}{2}\sigma_-\sigma_+\rho - \frac{1}{2}\rho\sigma_-\sigma_+\right).$$

[C] $t = \infty$:

Density matrix $\rho(\infty) = |\alpha|^2|0\rangle\langle 0| + |\beta|^2|1\rangle\langle 1|$.

4 Concluding Remarks

In this paper we tried to prove the Copenhagen interpretation of Quantum mechanics. Our method was based on embedding it into decoherence theory. Although the method is not complete it will become a starting point to give a complete proof to the Copenhagen interpretation.

In standard textbooks of QM decoherence theory is usually not contained, so it may be difficult to beginners (young students). See for example [10] or [11].

As far as we know this is the finest method at the present time.

References

[1] P. Dirac : **The Principles of Quantum Mechanics**, Fourth Edition, Oxford University Press, 1958.

[2] H. S. Green : **Matrix Mechanics**, P. Noordhoff Ltd, Groningen, 1965.

[3] Akio Hosoya : **Lectures on Quantum Computation** (in Japanese), SGC Library 4, Saiensu-sha Co., Ltd. Publishers (Tokyo), 1999.

[4] W. P. Schleich : **Quantum Optics in Phase Space**, WILEY–VCH, Berlin, 2001.

[5] K. Fujii : Superluminal Group Velocity of Neutrinos : Review, Development and Problems, Int. J. Geom. Methods Mod. Phys, **10** (2013), 1250083 (19 pages), arXiv:1203.6425 [physics].

[6] G. Lindblad : On the generator of quantum dynamical semigroups, Commun. Math. Phys, **48** (1976), 119.

[7] V. Gorini, A. Kossakowski and E. C. G. Sudarshan : Completely positive dynamical semigroups of N–level systems, J. Math. Phys, **17** (1976), 821.

[8] C. Zachos : Crib Notes on Campbell-Baker-Hausdorff expansions, unpublished, 1999, see http://www.hep.anl.gov/czachos/index.html.

[9] K. Fujii and et al : **Treasure Box of Mathematical Sciences** (in Japanese), Yuseisha, Tokyo, 2010.

I expect that the book will be translated into English.

[10] H. -P. Breuer and F. Petruccione : **The theory of open quantum systems**, Oxford University Press, New York, 2002.

[11] K. Hornberger : Introduction to Decoherence Theory, in “Theoretical Foundations of Quantum Information”, Lecture Notes in Physics, **768** (2009), 221-276, Springer, Berlin, quant-ph/061211.